



IJMARD 2014; 1(7): 16-18
www.allsubjectjournal.com
Received: 14-09-2014
Accepted: 18-11-2014
e-ISSN: 2349-4182
p-ISSN: 2349-5979

SK Saleem Bhasha
Research Scholar, University
College of Physical Education
and Sports Sciences, Acharya
Nagarjuna University,
Nagarjunanagar, Guntur,
India.

Y. Kishore
Dean, Faculty of Physical
Education and Sports Sciences,
Acharya Nagarjuna University,
Nagarjunanagar, Guntur,
India.

Correspondence:
SK Saleem Bhasha
Research Scholar, University
College of Physical Education
and Sports Sciences, Acharya
Nagarjuna University,
Nagarjunanagar, Guntur,
India.

Effect of aerobic training resistance training and concurrent training on resting heart rate among college boys

SK Saleem Bhasha, Y. Kishore

Abstract

The purpose of the study was to find out the effect of aerobic training, resistance training and concurrent training on resting heart rate among college boys. To achieve this purpose of the study, sixty college students were selected as subjects who were from the Acharya Nagarjuna University affiliated Colleges in Guntur. The selected subjects were aged between 18 to 22 years. They were divided into four equal groups of fifteen each, Group I underwent aerobic training, Group II underwent resistance training, Group III underwent concurrent training and Group IV acted as control that did not participate in any special training apart from their regular curricular activities. The subjects were tested on selected criterion variable such as resting heart rate prior to and immediately after the training period. The selected criterion variable such as resting heart rate was determined through using stop watch. The analysis of covariance (ANCOVA) was used to find out the significant differences if any, between the experimental group and control group on selected criterion variable. In all the cases, 0.05 level of confidence was fixed to test the significance, which was considered as an appropriate. The result of the present study has revealed that there was a significant difference among the experimental and control group on resting heart rate.

Keywords: resting heart rate -aerobic training-resistance training-concurrent training-sports.

1. Introduction

The primary objective of sports training is to stress various bodily systems to bring about positive adaptation in order to enhance sporting performance. To achieve this objective, coaches and athletes systematically apply a number of training principles including overload, specificity and progression, organized through what is commonly termed periodisation. The application of these principles involves the manipulation of various programme design variables including choice of exercise, order of training activities/exercises, training intensity (load and repetition), rest periods between sets and activities/exercises and training frequency and volume in order to provide periods of stimulus and recovery, with the successful balance of these factors resulting in positive adaptation. Aerobic exercise refers to exercise that involves or improve oxygen consumption by the body. Aerobic training increased cardio-respiratory endurance, which in turn increased VO_2 max, because of it increased level of hemoglobin. Resistance training is an integral part of an adult fitness program and of a sufficient intensity to enhance strength, muscular endurance and maintain fat free mass. Resistance training involves exercise in which the muscles exert a force against an external load. It is most commonly referred to as weight training. Such a training program should be individualized, progressive and specific in terms of the way muscles are likely to be used in the chosen sport. The physiological response to dynamic aerobic exercise is an increase in oxygen consumption and heart rate that parallels the intensity of the imposed activity and a curvilinear increase in stroke volume. The cardiovascular system, composed of the heart, blood vessels and blood responds predictably to the increased demands of exercise. With few exceptions, the cardiovascular response to exercise is directly proportional to the skeletal muscle oxygen demands for any given rate of work and oxygen uptake increases linearly with increasing rates of work. A person's maximum oxygen uptake is a function of cardiac output multiplied by the arterial-mixed venous oxygen difference. Cardiac output thus plays an important role in meeting the oxygen demands for work. As the rate of work increases, the cardiac output increases in a nearly linear manner to meet the increasing oxygen demand, but only up to the point where it reaches its maximal capacity. The resting heart rate can be obtained through auscultation, palpation or ECG recordings. When taking heart rate by auscultation, the bell of the stethoscope is placed to the left of the sternum, just above the level of the nipple. The heart beats can be counted.

1.1 Objectives of the study

The main objective of the study was to assess the effect of aerobic training, resistance training and concurrent training on resting heart rate which would help to enhance physical fitness of college boys. The present study was designed to obtain the data on the college boys from Acharya Nagarjuna University.

1.2 Statement of the problem

The purpose of the study was to determine the effect of aerobic training, resistance training and concurrent training on resting heart rate among college boys.

1.3 Delimitations

1. The study was delimited to Acharya Nagarjuna University affiliated colleges.
2. The study was delimited to 60 college students, their age was 18 to 22 years.
3. The study was restricted to the dependent variable is resting heart rate and independent variables are aerobic training, resistance training and concurrent training.

1.4 Significance of the Study

1. The findings of the study may be helpful for college students to apply aerobic, resistance and concurrent training which will help in better health and fitness.
2. The findings of the study would be helpful for the exercise physiologist to know the role of resting heart rate influence their physical fitness.
3. The results of the study may be helpful to fitness trainers, coaches, physical educationist and exercise physiologists to design proper training protocol for other populations.

2. Methodology

In the present study all the students studying in higher educational institutions of Acharya Nagarjuna University area were considered as population for the study. A representative sample of 60 college students in the age of 18-22 years was chosen as sample for the study. The selected participants were divided into four groups. Group I underwent aerobic training, group II underwent resistance training, group III underwent concurrent training and group IV act as control group. The experimental groups underwent eight weeks of training in their particular workout. For this study dependent variable is resting heart rate.

2.1 Test Administration – Resting Heart Rate

The subject was asked to sit and rest comfortably on a chair for 5 minutes. The pulse rate was determined at a radial artery. The investigator lightly pressed the index and middle finger against the radial artery in the groove on the lateral wrist. The pulse rate was counted for 30 seconds and multiplied by two to get the one minute value.

2.2 Analysis of Data

The data obtained were analyzed by analysis of covariance (ANCOVA). Analysis of covariance was computed for any number of experimental groups, the obtained 'F' ratio compared with critical F value for significance. When the F ratio was found to be significant, scheffe's post hoc test was used to find out the paired mean significant difference.

3. Results

3.1 Findings: The statistical analysis comparing the initial and final means of resting heart rate due to aerobic, resistance and concurrent training have been presented in Table I.

Table 1: Computation of Analysis of Covariation on Resting Heart Rate

Test	e.g. i	e.g. ii	e.g. iii	c.g.	f
Pre test	77.27	76.87	78.27	76.27	1.14
post test	72.07	73.00	73.33	75.33	3.05*
adjusted	71.98	73.27	72.34	76.15	31.12*

(The table value required for significance at .05 level with df 3 & 56 is 2.70 and 3 & 55 is 2.72).

Table I shows the analysed data of resting heart rate. The resting heart rate pre means were 77.27 for the aerobic training group, 76.87 for the resistance training group, 78.27 for concurrent training group and 76.27 for the control group. The resultant 'F' ratio of 1.14 was not significant at .05 levels indicating that the three groups were no significant variation. The post test means were 72.07 for the aerobic training group, 73.00 for the resistance training group, 73.33 for concurrent training group and 75.33 for the control group. The resultant 'F' ratio of 3.05 at .05 level indicating that was a significant difference. The difference between the adjusted post-test means of 71.98 for the aerobic training group, 73.27 for the resistance training group, 72.34 for concurrent training group and 76.15 for the control group yield on 'F' ratio 31.12 which was significant at .05 level.

The results of the study indicate that there is a significant difference among aerobic training, resistance training, concurrent training and control groups on the resting heart rate. To determine which of the paired means had a significant difference, Scheffe's post-hoc test was applied and the results are presented in Table II.

Table 2: Scheffe's Test for The Difference between the Adjusted Post-Test Paired Means of Resting Heart Rate

Adjusted Post-Test Means				Mean Diff.	Class Interval
Aerobic Training	Resistance Training	Concurrent Training	Control Group		
71.98	73.27			1.29	1.35
71.98		72.34		0.36	
71.98			76.15	4.17*	
	73.27	72.34		0.93	
	73.27		76.15	2.88*	
		72.34	76.15	3.81*	

The adjusted post test mean difference of resting heart rate between aerobic training and control group (4.17), resistance training and control group (2.88), concurrent training and control group (3.81) were greater than the required confidence interval of 1.35. The results of the study indicate that there were significant differences between control group and experimental groups and there was no significant difference between the experimental groups.

4. Discussion/Conclusions

The results of the study proved that there were significant differences between control group and aerobic training, resistance training and concurrent training group. The eight weeks of experimental treatment significantly influence on resting heart rate content in college students. However, there was no significant difference between experimental groups. The above results are supported by (Patricia and others, 2008, Wahab and Venkateswarul, 2010).

5. Recommendations

1. It was recommended that adequate steps may be taken to include aerobic, resistance and concurrent training in the physical education curriculum as these exercises significantly improves the resting heart rate of the subjects.
2. Similar study may be conducted on a larger population.
3. Similar study may be undertaken and its influence on psychological and biochemical parameters may be assessed.

6. References

1. Ansel Howard C, Prince Shelly J. Pharmaceutical Calculations: The Pharmacist's Handbook, LippincottWilliams & Wilkins, Philadelphia, USA, 2004.
2. Bompa TO. Theory and Methodology of Training, Dubuque, IA: Kendall/Hunt, 1983.
3. Frontera WR *et al.*, Strength Training and Determinants of VO₂ max in Older Women. *Journal of Applied Physiology* 1990; 68(1).
4. Pollock ML, Vincent KR. Resistance Training for Health: The President's Council on Physical Fitness and Sports Research Digest, December, Series 1996; 2:8.
5. Patricia CH Wong *et. al.*, Effects of 12 Week Exercise Training Programme on Aerobic Fitness, Body Composition, Blood Lipids and C-Reactive Protein in Adolescents with Obesity", *Ann Acad Med Singapore*, 2008, 37.
6. Wahab B Lawal, Venkateswarul Kankanala. Effects of Intermittent and Continuous Training on the Blood Pressure of Nigerian Adolescents and Children. *British Journal of Sports Medicine* 2010; 44:28.
7. Clarke H Harrison, Clarke David H. *Advanced Statistics*, New Jersey: Prentice Hall Inc, 1972.