



International Journal of Multidisciplinary Research and Development



IJMRD 2015; 2(1): 216-220
www.allsubjectjournal.com
Received: 15-12-2014
Accepted: 08-01-2015
e-ISSN: 2349-4182
p-ISSN: 2349-5979
Impact factor: 3.762

Sri Ranjani Sivapalan
Senior Lecturer,
Gr 1 Unit of Siddha Medicine,
University of Jaffna,
Jaffna 40000, Sri Lanka.

Phytochemical study on medicinal plant – *Sida cordifolia* Linn

Sri Ranjani Sivapalan

Abstract

Texts state that the *Sida cordifolia* has analgesic, antispasmodic, anti-inflammatory, hypoglycemic and hepatoprotective properties. This species is found in tropical and subtropical regions of Sri Lanka and India. The aim of this study is to determine the phytochemical constituents of *Sida cordifolia*. phytochemistry helps in standardizing the herbal preparations and possibly relate the constituents to their medicinal/pharmacological uses. Dried material was grounded to coarse powder and stored in airtight container. It was then extracted with ethanol. The dry powder of sample was observed under U.V. light to evaluate the fluorescence. Chemical tests were performed on ethanol extract. The quantitative studies revealed that *Sida cordifolia* possessed alkaloids (1.99 mg/kg), flavonoids (0.92 mg/kg), lignin (0.08 mg/kg), glycosides (0.19 mg/kg), saponins (0.17 mg/kg), phytosterols (0.02 mg/kg), fixed oils (0.18 µg/lit). Phytochemicals act in numerous ways to assist the human body in combating disease and health problems. The preliminary phytochemical screening tests may be useful in the detection of the bioactive principles and subsequently may lead to the drug discovery and development. It is also justify the traditional medical uses and the claims about the therapeutic values of this plant as curative agent. Further to this the isolation, identification, characterization and elucidation of the structure of the bioactive compounds of *Sida cordifolia* would be obtained with a view to obtain useful chemotherapeutic agent.

Keywords: Phytochemical, *Sida cordifolia*, Medicinal plants, *Thuththi*, *Bala*, *Babila*

1. Introduction

Plant and plant extracts have formed important position in modern medicine, due to their chemical and medicinal contents found in the natural form. The secondary metabolites represent a large reservoir of structural moieties which work together exhibiting a wide range of biological activities [1]. Medicinal plants form the major source of drugs in all the traditional systems of medicine practiced in Sri Lanka, India viz., Siddha, Ayurveda, Unani and Homeopathy. There is a growing importance in medicinal plants and traditional health systems providing health care for a wider population across the globe, especially, in the developing countries. The World Health Organization (WHO) currently encourages, recommends and promotes traditional remedies in health care programs as they are easily available at low cost, comparatively safe and are culturally acceptable [2]. Every culture has depended on the healing power of herbs.

Traditional medical texts states that the *Sida cordifolia* posses analgesic, antispasmodic anti-inflammatory, hypoglycemic and hepato-protective activities [3, 4]. This plant is used in bronchial asthma, nasal congestion, skin disease, urinary diseases, obesity, cardiac diseases, bleeding haemorrhoids and preparation of analgesic massage oils [5]. Kirtikar and Basu stated that *Sida cordifolia* is used in arrhythmia, hemiplegia, sciatica, neuritis, neuralgia, epilepsy, rheumatism, anorexia, fatigue, impotence, spermatorrhea, gonorrhea, cystitis, leucorrhea, urinary frequency, diabetes, diarrhoea, dysentery, hemorrhoids, chronic fever [6]. This species is found in tropical and subtropical regions of Sri Lanka and India. It grows as wasteland weed. It is also known as the “Bala” in Hindi and Sanskrit [7]. The aim of this study is to determine the phytochemical constituents of *Sida cordifolia* and possibly relate the constituents to their medicinal/pharmacological uses. Phytochemical studies of the plant are necessary for standardization, which helps in understanding the significance of phytoconstituents in terms of observed activities.

Correspondence:
Dr. Sri Ranjani Sivapalan
Senior Lecturer,
Gr 1 Unit of Siddha Medicine,
University of Jaffna,
Jaffna 40000, Sri Lanka.

Phytochemistry also helps in standardizing the herbal preparations.

Sida cordifolia Linn (Figure 1.1) is belonging to the family Malvaceae. Found throughout the tropical and sub-tropical plains of India and Sri Lanka.



Kingdom	Plantae – Plants
Subkingdom	Tracheobionta – Vascular plants
Superdivision	Spermatophyta – Seed plants
Division	Magnoliophyta – Flowering plants
Class	Magnoliopsida – Dicotyledons
Subclass	Dilleniidae
Order	Malvales
Family	Malvaceae – Mallow family
Genus	<i>Sida</i> L. – fanpetals
Species	<i>Sida cordifolia</i> L.

Fig 1.1: *Sida cordifolia*

English: Country mallow,

Sanskrit: Bala

Tamil: Thuththi

Sinhala: Bala, babila

1.1 Botanical Description

Plant- an erect, perennial shrub, upto 1m. Tall. Stem- Ascending, terete or sulcate, softly villous and densely stellate-pubescent all over. Leaves- Ovate or ovate-oblong, obtuse or subacute at apex. Flowers- Yellow, peduncles, axillary, jointed much above the panicles, upper flowers nearly sessile and fasciculate towards the tip of the branches forming subsapicate inflorescence. Fruits- Subdiscoid, 6-8 mm across, mericarps 10, 3 sided. Seeds- Trigonous, glabours, tufted-pubescent near the hilum. Flowering & Fruiting Time - August-November in Jaffna conditions.

2. Materials and Methods

2.1 Collection and Identification of plant materials

Areal parts of *Sida cordifolia* for the proposed study were collected during the month of July-August 2012 from Jaffna district, Sri Lanka and care was taken to select healthy plants. The identity of the plant specimens was confirmed by the use of local Floras [8]. The botanical identify was authenticated by a botanist, Department of Botany,

University of Jaffna. The collected plant materials were washed thoroughly with tap water and dried under shade for ten days. Dried material was grounded to coarse powder and stored in airtight container. It was then extracted with ethanol.

The dry powder of sample was observed under U.V. light to evaluate the fluorescence. Chemical tests were performed on ethanol extract.

2.2 Qualitative analysis - Phytochemical study

- **Preparation of ethanol extract:** About 600 gm of powered material was taken in a clean, flat-bottomed glass container and soaked in 800 ml of 95% ethanol. The container with its contents was sealed and kept for a period of 7 days accompanying occasional shaking and stirring. The whole mixture then underwent a coarse filtration by a piece of clean, white cotton material. Then it was filtered through whatman filter paper. The filtrate thus obtained was concentrated using a rotary evaporator to get the extract.
- **Phytochemical Screening:** Phytochemical screening means to investigate the plant material in terms of its active constituents. The plant extract was subjected to qualitative tests for the identification of the phyto-constituents present in it viz, alkaloids, carbohydrates, glycosides, phytosterols, fixed oils & fats, phenolic compounds & tannins, proteins and free amino acids, flavonoids, lignins and saponins [9, 10, 11, 12].
- **Test for Phenolic compounds:** Two to three drops of 1% ferric chloride (FeCl_3) solution was added into 2ml of extract. Phenolic compounds produce a deep violet colour with ferric ions.
- **Test for Flavonoids - Shinoda Test:** To the alcoholic solution of the extract a few fragments of magnesium ribbon were added. To this concentrated hydrochloric acid (HCl) was added drop wise. Magenta colour was produced after few minutes which are the characteristic reaction of flavonoid.
- **Test for Tannins -Ferric Chloride test:** Water extract was treated with 15 % ferric chloride test solution. The resultant colour was noted. A blue colour indicates condensed tannins; a green colour indicated hydrolysable tannins.
- **Test for Saponins:** The extract was diluted with 20ml of distilled water and it was agitated on a graduated cylinder for 15 min. the presence of saponins was indicated by the formation of 1cm layer of foam.
- **Test for Alkaloids - Mayer's test:** The Extract was dissolved in chloroform. The chloroform was evaporated and the residue was acidified and added few drops of Mayer's reagent (Potassium Mercuric Iodide). Alkaloids are precipitated by Mayer's reagent to give a cream coloured precipitate.
- **Wagner's Test:** The extract was dissolved in chloroform. The chloroform layer was evaporated, to the residue was acidified and added few drops of Wagner's reagent (Iodine in Potassium Iodide). Orange precipitate indicates alkaloids.

- **Test for the carbohydrate:** The aqueous extract 5 ml was treated with the reagent of the starch (iodine). Any shift to blue violet indicates the presence of starch.
- **Test for Glycosides - Fehling's Test for reducing sugars (In Glycosides):** The extract was re-dissolved in water on the water bath. To 2 ml of the solution, in the test tube was added, 1ml each of Fehling's solutions A and B. The mixture was shaken and heated in a water bath for 10min. The colour obtained was recorded. A brick-red precipitate indicates reducing sugar.
- **Test for proteins - Xanthoproteic Test:** Extract was treated with few drops of Concentrated HNO_3 . Formation of yellow colour indicates the presence of proteins.
- **Test for Phytosterols - Liebermann-Burchard's Test:** One gram of the extract was treated with chloroform and filtered. The filtrate was treated with few drops of acetic anhydride boiled and cooled, concentrated sulphuric acid was added through the sides of the test tube. The formation of brown coloured ring at the junction of two liquids confirmed the presence of steroids.
- **Test for fixed oil and fats:** Press the extract in between the two filter papers, a permanent stain indicates the presence of fixed oil. Extract was treated with few drops of 0.5N potassium hydroxide and few drops of phenolphthalein and heat, formation of soap indicates the presence of fixed oil and fats.

2.3 Fluorescence Analysis

Fluorescence analysis of the plant powder was observed in daylight and UV light (254nm) in a UV chamber ^[13, 14] after treating with different chemical reagents is reported (Table 3.1). It can be as a diagnostic tool for testing the adulterations.

2.4 Quantitative study of Phytochemicals

- **Determination of Alkaloid ^[15]:** 5 g of the sample was weighed into a 250 ml beaker and 200 ml of 10% acetic acid in ethanol was added and covered and allowed to stand for 4h. This was filtered and the extract was concentrated on a water bath to one-quarter of the original volume. Concentrated ammonium hydroxide was added drop wise to the extract until the precipitation was complete. The whole solution was allowed to settle and the precipitated was collected and washed with dilute ammonium hydroxide and then filtered. The residue is the alkaloid, which was dried and weighed.
- **Determination of Flavonoid ^[16]:** 10 g of the plant sample was extracted repeatedly with 100 ml of 80% aqueous methanol at room temperature. The whole solution was filtered through whatman filter paper No 42 (125 mm). The filtrate was later transferred into a crucible and evaporated into dryness over a water bath and weighed to a constant weight.
- **Determination of Saponin ^[17]:** 20 g of each grounded sample was put into a conical flask and 100cm³ of 20% aqueous ethanol was added. Then the flask was heated on a hot water bath for 4 h. with constant stirring at about 55°C. The mixture was then filtered and the

residue was again extracted with another 200 ml 20% ethanol. The combined extract was reduced to 40 ml on a hot water bath at about 90 °C. The concentrate was transferred into a 250 ml separatory funnel, added 20 ml diethyl ether in it followed by vigorous shaking. The aqueous layer was recovered while the ether layer was discarded. The purification process was repeated. 60 ml of n-butanol was added. The combined n-butanol extracts were washed twice with 10 ml of 5% aqueous sodium chloride. The remaining solution was heated in a water bath. After evaporation the samples were dried in oven, weighed and saponin content was calculated as percentage.

➤ Determination of Glycosides

Test 1: Extract of 200mg of the sample was taken warmed in a test tube with 5ml of diluted 10% sulphuric acid on the water bath at 100°C for 2 minutes. Centrifuged; pipette off the supernatant. The acid extract was neutralized with 5% solution of NaOH. Added 0.1 ml of Fehling's solution A and then Fehling's solution B until alkaline (tested with pH paper) and heated on the water bath for 2 minutes. Noted the quality of red precipitate formed and compared with that formed in test 2.

Test 2: Extract of 200mg of the sample was taken an added 5ml of water instead of Sulphuric acid. After boiling added a volume of water equivalent to the volume of NaOH used in the test 1, step 2 and above. Added 0.1 of Fehling's solution B until alkaline (tested with pH paper) and heated on the water bath for 2 minutes. Note the quantity of red precipitate formed (test 2). Compared the quality of the precipitate formed in test 2 that formed in test 1. The precipitate in test 1 was greater than that in test 2. That indicated the presence of glycoside in the crude drug ^[18].

- **Determination of Lignins:** 2gms of the oven-dry material passed through an 80-mesh sieve were extracted in a Soxhlet apparatus with 200ml, 95% alcohol for 4 hours. The extracted residue was transferred to a round-bottomed flask of 300ml capacity and extracted with 150ml boiling water under a reflux for 1 hour. The contents of the flask are then filtered and residue was transferred back to the flask and hydrolyzed with 150ml. 5% H one hour. The hydrolyzed product was then collected in a weighed sintered glass crucible after being washed free of acid. The residue is weighed and well powdered. 0-2g lots were weighed out into 1000 ml beakers and treated with 20ml 72% H_2SO_4 . The powder was well mixed with the acid and allowed to stand overnight, at a temperature varying between 20 and 220 (this was the range between 4p.m and 8a.m the following morning). The following morning the contents of the beakers were made up to 800ml. with water and boiled for 2 hours, the volume being maintained by occasional addition of water. Leave the beakers overnight to settle down the Lignin's, thereby facilitating filtration. The filtration was done under

suction in Gooch crucibles with No. 42 Whatmann filter paper. After drying, the precipitates were weighed and ignited and the lignin's calculated on an ash-free basis.

- **Determination of Phytosteroids:** 1g of powdered dried sample was extracted 3 times using a vortex mixture (15min) with 7.5 ml chloroform. All the extracts were combined and evaporated to dryness. This chloroform extract contains free sterols and terpenoids. The residue was hydrolysed with 2N HCl in methanol (2hr, 75-80° c), neutralized with 10N NaOH, then diluted with 25ml water, and then the steroidal alkaloids and sapogenins were extracted 3 times with 10ml chloroform. The chloroform phase was collected and evaporated to dryness. The amounts of phytosteroids were weighed.
- **Determination of Fixed Oil:** Transfer a 50gm of the air dried, crushed drug to an extraction thimble, extract with Solvent ether in a continuous extraction apparatus (Soxhlet extractor) for 6 hours. Filter the extract quantitatively into a tared evaporating dish and

evaporate off the solvent on a water bath. Dry the residue at 105° to constant weight. Calculate the percentage of ether-soluble extractive with reference to the air-dried drug.

3. Results and Discussion

The result of Fluorescence analysis, Qualitative and quantitative analysis of *Sida cordifolia* have been presented and discussed herewith.

When physical and chemical parameters are inadequate as it often happens with the powdered drugs, the plant material may be identified from their adulterants on basis of fluorescence study^[19].

Fluorescence analysis of drug showed in table 3.1 is an important parameter in detecting adulteration or improper handling of drugs. It can serve as a valuable source of information and provide appropriate standards to establish the quality of this plant material in future study or application.

Table 3.1: Fluorescence analysis of *Sida cordifolia*

Serial No	Chemical test	<i>Sida cordifolia</i> Day light	<i>Sida cordifolia</i> UV light
1.	Sample	Yellow	Dark Yellow
2.	Powder + 1N aq. NaOH	Yellow	Green
3.	Powder+ 1N alc. NaOH	Yellowish green	Dark green
4.	Powder + 1N HCl	Greenish yellow	Green
5.	Powder + 50% HNO ₃	Light green	Dark green
6.	Powder + 50% H ₂ SO ₄	light yellow	Green
7.	Powder + Methanol	Green	Blackish green
8.	Powder + NH ₃	White	Green
9.	Powder + I ₂	Reddish orange	Yellowish brown
10.	Powder + FeCl ₃	Brownish orange	Dark brown

Table 3.2: Phytochemical Screening of the Ethanolic extract of *Sida cordifolia*

Serial No	Constituents	<i>Sida cordifolia</i> ethanolic extract
1.	Alkaloids	+
2.	Carbohydrates	+
3.	Glycosides	+
4.	Saponins	+
5.	Tannins	-
6.	Phytosterols	+
7.	Proteins	+
8.	Flavonoids	+
9.	Lignin	+
10.	Volatile oils	-
11.	Fixed oil and fats	+

Result from the phytochemical screening test in Table 3.2 indicated the presence of phenolic compounds, flavanoid, phytosterol, alkaloids, glycoside and saponin in the aerial parts of the *Sida cordifolia*. The phenolic compounds are considered as being a major group to the number of the secondary metabolites that contributes to the antioxidant activity of the plant. The presence of phenolic compounds in the plant indicates that this plant may have the ability as an anti-microbial agent^[20].

It has been reported that most active principles in plants are frequently flavonoids, steroids, glycosides and alkaloids. These phytoconstituents may be responsible for the many pharmacological actions of the plant like wound healing^[21] cholesterol lowering^[22] and antidiabetic activity. Phytosterols are an important breakthrough in the human fight against high cholesterol. It has been known that plant steroids, flavonoids are antioxidants. These antioxidants are compounds that reduce the formation of free radicals or react with and neutralize them thus potentially protecting the cell from oxidative damage^[23].

Table 3.3: Quantitative phytochemical evaluation of *Sida cordifolia*

Serial No	Parameters	<i>Sida cordifolia</i>
1	Total Alkaloids (mg/kg)	1.99
2	Glycosides (mg/kg)	0.19
3	Saponins (mg/kg)	0.17
4	Phytosterols (mg/kg)	0.02
5	Total Flavonoids (mg/kg)	0.92
6	Lignin (mg/kg)	0.08
7	Fixed oil (µg/lit)	0.18

Table 3.4: Results of the Macronutrient analysis of *Sida Cordifolia*

Serial No	Constituents	<i>Sida cordifolia</i> ethanolic extract
1	Carbohydrates	+
2	Reducing Sugar	+
3	Protein	+
4	Fixed oil and fats	+

The results of the micronutrient analysis showed the presence of carbohydrate, fats, protein, reducing sugar and oil were present (Table 3.4).

On the other hand, the macronutrients; proteins, carbohydrate and reducing sugar are involved in the energy giving and body building function of *Sida cordifolia*.

4. Conclusion

The results obtained from the phytochemical analysis of the areal part of *Sida cordifolia* showed the presence of alkaloids, steroids, flavonoids, Phytosterols, lignins, and macronutrient analysis revealed the presence of proteins, carbohydrates, reducing sugar, fats and oil. This study justifies the use of *Sida cordifolia* in the treatment of many ailments like diabetic mellitus, carcinoma, heart disease, asthma. Non-nutrient (phytochemicals) content which act as antioxidant against dangerous free radicals in the body system. In this study suggest the identified phytochemical compounds may be the bioactive constituents which are medicinally valuable. Therefore, extracts from these could be seen as a good source for useful drugs and it is suggested that further work is in progress to isolate, purify, and characterize the active constituents responsible for the activity of these plants.

5. References

- Narendra Devanaboyina N, Ram Sharmila BG, Kumar G, Rajasekhara PM. Cholesterol-lowering activity of the aqueous fruit extract of *Trichosanthes dioica* Roxb. In normal and streptozotocin diabetic rats. *J Clin Dia Res* 2007; 1(4):561-569.
- World Health Organization, Organization Mondiale De La Sante, Quality control methods for plant materials, 559, Rev,1, Original English, 1998, 08-67.
- Anilkumar M. Ethnomedicinal plants as antiinflammatory and analgesic agents. Ed. Debprasad Chattopadhyay. *Ethnomedicine: A Source of Complementary Therapeutics*, 2010, 267-293.
- The Indian Pharmacopoeia., Govt. of India, Ministry of Health and Family Welfare, The controller of Publication, 1996, 53, 54, 89.
- Nadkarni's KM. *Indian Materia Medica*. Vol. Popular Prakashan, Bombay, 1976, 1137.
- Kirtikar KR, Basu BD. *Indian Medicinal Plants*. 2nd Edn. Bishen Singh Mahendra Pal Singh, Dehradun, 1980, 312. Nayar, Chopra. *Glossary Indian med: plants*, PID. New Delhi, 1956, 227.
- Kirtikar KR, Basu BD. In: Blatter E, Caius JF, Mhaskar KS (Eds.), *Indian Medicinal. Plants*. Lalit Mohan Basu, Allahabad, India, 1933, 2390–2392.
- Kokate CK. *Practical Pharmacognosy*. Vallabh Prakashan, New Delhi, 1994, 107-113.
- Harborne JB. *Phytochemical Methods*. Chapman & Hall, London, 1998, 60-66.
- Sofowora A. *Medicinal Plants and Traditional Medicine in Africa*. John Wiley and Sons Ltd. Chichester, 1982, 142-145.
- Trease GE, Evans WC. *Pharmacognosy* 2nd Edn. Braille Tirideland Macmillan Publishers, 1989.
- Chase CR, Pratt R. Fluorescence of powdered vegetables drugs with particular reference to development of a system of identification. *J.Am Pharma Assoc* 1949; 38:324-31.
- Kokoshi CJ, Kokski RJ, Sloma FJ. Methods for the analysis of various compound in plants. *Journal of Pharmacognostical Association*, 1958, 10-716.
- Harborne JB. *Methods of plant analysis In Phytochemical Methods*. Chapman and Hall, London, 1973.
- Bohm BA, Kocipai Abyazan R. Flavonoid and condensed tannins from the leaves of *Vaccinum raticulation* and *Vaccinum calcyimium*. *Pacific Sci* 1994; 48:458-463.
- Obadoni BO, Ochuko PO. Phytochemical studies and comparative efficacy of the crude extract of some homeostatic plants in Edo and Delta states of Nigeria". *Global J Pure Appl Sci* 2001; 8:203-208.
- Zafar R, Lalwani M. *Tribulus terrestris* Linn. A review of current knowledge", *Indian Drugs* 1989;27(3):148-158.
- Rai MK, Shende S, Strasser RJ. JIP test for fast fluorescence transients as a rapid and sensitive technique in assessing the effectiveness of arbuscular mycorrhizal fungi in *Zea mays*: Analysis of chlorophyll a fluorescence. *Plant Biosystems* 2008; 142(2):191-198.
- Harisharanraj R *et al.* Evaluation of the Chemical Composition *Rauwolfia serpentina* and *Ephedra vulgaris*. *Advances in Biological Research* 2009; 3(5-6):174-178.
- Shivhare Y, Singour PK, Patil UK, Pawar RS. Wound healing potential of the methanolic extract of *Trichosanthes dioica* Roxb (fruits) in rats. *J Ethnopharmacol* 2010; 127:614-619.
- Shrmila BG, Kumar G, Rajasekhara PM. Cholesterol-lowering activity of the aqueous fruit extract of *Trichosanthes dioica* Roxb. In normal and streptozotocin diabetic rats". *J Clin Dia Res* 2007; 1(4):561-569.
- Delanty N, Dichter MA. Antioxidant therapy in neurologic disease., *Arch Neurol*, Department of I.S.M. & H., New Delhi, Part-I, Volume -I, 2 Appendix 2000; 57(9):1265-70.