



Volume: 2, Issue: 8, 190-192
Aug 2015
www.allsubjectjournal.com
e-ISSN: 2349-4182
p-ISSN: 2349-5979
Impact Factor: 3.762

Nilesh G Patel
Final Year Resident,
Community Medicine
Department, B. J. Medical
College, Ahmedabad,
Gujarat, India.

Parimal Patel
Final Year Resident,
Community Medicine
Department, B. J. Medical
College, Ahmedabad,
Gujarat, India.

Mitin Parmar
Final Year Resident,
Community Medicine
Department, B. J. Medical
College, Ahmedabad,
Gujarat, India.

MK Lala
Associate Professor,
Community Medicine
Department, B. J. Medical
College, Ahmedabad,
Gujarat, India.

Correspondence
Nilesh G Patel
Final Year Resident,
Community Medicine
Department, B. J. Medical
College, Ahmedabad,
Gujarat, India.

A Comparative study on Pre-hypertension and its determinants in first year and final year students of B.J. Medical Collage, Ahmedabad

Nilesh G Patel, Parimal Patel, Mitin Parmar, MK Lala

Abstract

Pre-hypertension is associated with an increased risk of the development of hypertension and subsequent cardiovascular disease and raises mortality risk. Pre-hypertension is emerging health problem in adolescents now a day. The aim of this study was to determine the prevalence of pre-hypertension, to explore the association between pre-hypertension and established cardiovascular risk factors and compare them among a first year and final year students of B.J.M.C Ahmedabad. In this cross-sectional study a representative sample of 200 participants was selected using a Random sampling method. Pre-hypertension was observed among 73 out of Total 200 students, 31 from first year and 42 final year student, 40 for Boys and 33 for Girls. The pre-hypertensive group had higher body mass index, higher percentage of smoking, ate more added salt and less physical activity than did the normotensive group. Prevalence of pre-hypertension was higher among final year students than first year students due to heavy study stress. Therefore, primary prevention strategies should concentrate on reducing overweight and obesity. Diet education should be given to overcome this problem. Stress is another major predictor in medical students so Medical authority should make some strategies to relieve stress in medical students. Keywords: Pre-hypertension, Cardiovascular risk factor, Obesity, BMI, stress factor, Medical students.

Keywords: Comparative study, Pre-hypertension, students, B.J. Medical Collage

Introduction

The global burden of non-communicable diseases is emerging as a major public health challenge in the world.¹ One thing is clear: "when it comes to NCDs, inaction is not an option."² According to WHO, chronic disease is modern epidemic. 60% of all deaths and 47% of burden of diseases are due to non-communicable diseases; these figures are expected to rise to 73% and 60%, respectively, by 2020.³

Hypertension is a well-known risk factor for the development of CVD, heart attack, and stroke.³ Hypertension is responsible for 57% of stroke deaths and 24% of coronary heart disease deaths in India.⁴ Due to change in life style, environmental and Socio-demographic risk factors the prevalence of Pre-hypertension and hypertension is increasing constantly.

Pre-hypertension is associated with an increased risk of the development of hypertension and subsequent cardiovascular disease and raises mortality risk. Persons with pre-hypertension have a greater risk of developing hypertension than do those with lower blood pressure levels.⁵

Pre-hypertension is also an emerging health problem in adults and adolescents now a day. Medical Students are more prone to be exposed with established cardiovascular risk factors.

Aims

The study was aimed to know the prevalence of Pre-hypertension and its determinants in First year and Final year students of B.J.M.C Ahmedabad and compare them.

Objectives

To determine the prevalence of pre-hypertension

To explore the association between pre-hypertension and established cardiovascular risk factors
Compare those risk factors among a first year and final year students of B.J.M.C Ahmedabad.

Methodology

Sample Size: Prevalence of pre-hypertension was 34% in various studies among adolescents and adults. Sample size was calculated by the Following formula.

Sample Size (N) = 4pq/L2
 Where, p = Prevalence of pre-hypertension=34%
 q = 100-p = 66%
 L = 20% of p = 6.8
 So Sample size (N) = 4*34*66/46.51

The estimated sample size was 193. Sample size of 200 was decided. 100 students from first year and 100 students from final year.

Study design: In this cross-sectional study, Participants were selected using a Random sampling method. (Chit method)

Study Area: B. J. Medical College, Ahmedabad

Study Population: First year and Final year students of B. J. M. C. Ahmedabad.

Data collection: All Students were interviewed with pre-designed questionnaire.

Blood pressure was measured after the students had rested for at least 5 minutes and from right arm placed at the heart level. Two measurements were taken by a mercury sphygmomanometer with at least 5 minutes between successive measurements and lower one was recorded.

According to the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and treatment of High Blood Pressure (JNC 7) in 2003, prehypertension is defined as blood pressure range with a systolic BP of 120–139 millimeters of mercury (mmHg) and/or a diastolic BP of 80–89 mmHg.

Body weight was measured with light cloth and without shoes, a common weighing machine was used for that.

Height was measured to the nearest 0.5 cm, without shoes using a measuring tape

Body mass index (BMI) was calculated as weight (in kilograms) divided by height (in meters) squared.

Data Analysis: Statistical analysis was performed using Microsoft excel and EPI INFO version 7 software.

Results

Pre-hypertension was observed among 73 (36.5%) out of 200 students, 34 (34%) from first year and 42 (42%) final year student while 5(5%) were hypertensive in first year and 8(8%) in final year.

Table 1: Presence of different risk factors among Normotensive and Pre-hypertensive FIRST YEAR Students

Risk Factors	Normotensive (61) (61%)	Pre-hypertensive (34) (34%)	P value
Family history of hypertension	11 (18.32%)	6 (17.64%)	0.9643
Stress Factor	10 (16.39%)	8 (23.52%)	0.3948
Male sex	37 (60.65%)	18 (52.94%)	0.4653
Excessive salt intake	30 (45.45%)	19 (55.88%)	0.5307
Smoking	14 (21.21%)	9 (26.47%)	0.7014
Lack of physical activity	24 (36.36%)	16 (47.05%)	0.4653
Overweight/obese	32 (48.48%)	19 (55.88%)	0.7482

Year	Normal	Overweight	Obese	Total Pre-hypertensive	P Value
First Year	15(41.18%)	16 (47.05%)	3(11.77%)	34 (100%)	0.7452
Final Year	16(38.10%)	19 (45.23%)	7(16.67%)	42 (100%)	0.08

Discussion

The pre-hypertensive group had higher body mass index, higher percentage of smoking, ate more added salt and less physical activity than did the normotensive group. (Table: 1)

Table 1 showing different risk present among first year students of B.J.M.C. Male sex, excessive salt intake and overweight and obese were major risk factors which were observed in first year students of B.J.M.C, Ahmedabad. While Table 2 showing different risk present among final year students of B.J.M.C. Major risk factors which were observe in final year students were almost same as in first year students of B.J.M.C, Ahmedabad.

Table 2: Presence of different risk factors among Normotensive and Pre-hypertensive FINAL YEAR Students

Risk Factors	Normotensive (61) (61%)	Pre-hypertensive (34) (34%)	P value
Family history of hypertension	12 (24%)	10 (23.80%)	0.8301
Stress Factor	9 (18%)	17 (40.47%)	0.017
Male sex	27 (54%)	24 (57.14%)	0.7629
Excessive salt intake	21 (42%)	22 (52.38%)	0.3202
Smoking	11 (22%)	11 (26.19%)	0.6390
Lack of physical activity	19 (38%)	18 (42.85%)	0.6360
Overweight/obese	22 (44%)	26 (61.90%)	0.08

Table 3: Association of Pre-hypertension with Stress

Year	Total Pre-hypertensive Students	Pre-hypertensive Students With Stress
First Year	34	8(23.52%)
Final Year	42	17(40.47%)

First Year: chi square value 0.401, df =1, P=0.52657 (P>0.05), Test is not significant.

Final Year: chi square value 5.591, df =1, P=0.018059 (p<0.05), Test is not significant.




Fig 1: Distribution of Pre-hypertensive according to BMI

(Table: 2) These findings are consonant with previous reports on risk factors in pre-hypertension for cardiovascular disease.⁸ Prevalence of pre-hypertension was higher among final year students than first year students may be due to heavy study

stress. Analysis showed that Stress is major risk factor to developing Pre-hypertension and also hypertension. (Table: 4) Analysis showed that obesity and overweight were the strongest predictors of pre-hypertension.

Conclusions and Recommendation

Overweight and obesity are major determinants of the high prevalence rate of pre-hypertension detected in students of B.J.M.C Ahmedabad. Therefore, primary prevention strategies should concentrate on reducing overweight and obesity. Diet education should be given to overcome this problem.

Stress is another major predictor in Final Year medical students so Medical authority should make some strategies to relieve stress in medical students.

The normotensive Group also having higher percentage of these risk factors so they are also prone to develop pre hypertension in future.so preventive step should take in Normotensive students like health education.

Reference

1. Murray CJL, Lopez AD. The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 projected to 2020. Cambridge (MA): Harvard School of Public Health; 1996.
2. By: David E. Bloom and Elizabeth T. Cafiero Nov 7th 2012. (Available from: URL: <http://forumblog.org/2012/11/are-non-communicable-diseases-taking-aim-at-india/>) (Last accessed 2013 on Dec 4.)
3. World Health Report. Geneva: WHO, 2002
4. R. Trends in hypertension epidemiology in India. Journal of Human Hypertension 2004; 18:73–78
5. Greenlund KJ, Croft JB, Mensah GA. Prevalence of heart disease and stroke risk factors in persons with prehypertension in the United States, 1999-2000. Arch Intern Med 2004; 164: 2113-8.
6. Daar AS, Singer PA, Persad DL, Pramming SK, Matthews DR, Beaglehole R, *et al*. Grand challenges in chronic non-communicable diseases. Nature. 2007 Nov 22;450(7169):494-6
7. Epidemiology of Hypertension Who, Global11, 12 © SUPPLEMENT TO JAPI • FEBRUARY 2013 • VOL. 61
8. Kshirsagar AV, Carpenter M, Bang H, Wyatt SB, Colindres RE. Blood pressure usually considered normal is associated with an elevated risk of cardiovascular disease. Am J Med. 2006; 119:133–141. [PubMed: 16443415]