



Analyzing the status of genetic enhancements in mung bean

Emanuel Hussein, Joseph M Matee

Department of Food Science and Technology, Mbeya University of Science and Technology, Mbeya, Tanzania

Abstract

Mung bean (*Vigna radiata*), a vital legume crop, has long been celebrated for its nutritional value and adaptability to diverse agricultural environments. However, global challenges such as climate change, population growth, and the increasing need for sustainable farming practices necessitate advanced genetic enhancements to improve its resilience, productivity, and nutritional quality. Despite significant strides in breeding and biotechnological approaches, there remains a crucial need to systematically review and analyze the status of genetic enhancements in mung bean. This review aims to collate recent advancements in genetic improvements, assess the integration of traditional and modern techniques, and identify future research directions that could further bolster the crop's contribution to food security and agricultural sustainability.

Keywords: Mung bean (*Vigna radiata*), genetic enhancements, agricultural environments

Introduction

Mung bean serves as a cornerstone in the diets of millions worldwide, providing essential proteins, vitamins, and minerals. It also plays a crucial role in sustainable agriculture due to its nitrogen-fixing ability, enhancing soil health and reducing the need for synthetic fertilizers. However, the full potential of mung bean is yet to be unlocked, as its cultivation is hampered by various biotic and abiotic stresses, including pests, diseases, drought, and salinity. The escalation of these challenges, driven by changing climatic conditions and the ever-growing global population, underscores the urgent need for enhanced genetic interventions. Historically, the improvement of mung bean was largely reliant on conventional breeding techniques. While successful in developing strains with desirable traits, the pace and scope of these improvements are insufficient to meet current and future demands. The advent of molecular biology and biotechnology offers new avenues for the genetic enhancement of mung bean, promising rapid, precise, and wide-ranging improvements beyond what traditional breeding can achieve. These technologies enable the direct manipulation of the crop's genetic material, offering solutions to enhance yield, stress tolerance, and nutritional value effectively.

Main Objective

The main objective of this review is to provide a comprehensive overview of the current status of genetic enhancements in mung bean.

Literature Review

(Ramanujam, Tiwari, & Mehra, 2004) ^[1] have found substantial genetic divergence among mung bean populations, indicating significant potential for breeding

high-performance hybrids. Characteristics such as flowering time, maturity, seed density, and seed size have been identified as major contributors to this divergence, which could be exploited for enhancing yield and other desirable traits.

(Chen *et al.*, 2015) ^[2] developed and utilized microsatellite SSR markers to assess genetic diversity and population structure among mung bean germplasms. This research provided insights into the low polymorphism within the species and identified distinct subgroups within the cultivated and wild mung bean accessions, suggesting pathways for further genetic improvement.

(Chen *et al.*, 2015) ^[3], Transcriptome sequencing has been used to identify potential molecular markers in mung bean, aiding in the construction of genetic linkage maps and facilitating the genetic mapping of traits. This approach has led to the identification of a large number of EST-SSR markers, which are valuable for breeding and genetic studies.

(Ha & Lee, 2019) ^[4], Efforts have been made to mine useful alleles from diverse germplasms for mung bean breeding. With the availability of the mung bean genome sequence, there's a push towards utilizing genomic information to overcome the challenges in mung bean breeding, such as resistance to biotic and abiotic stresses and improvement of nutritional quality.

(Dahiya *et al.*, 2015) ^[5], Mung bean is recognized for its high nutritional and economic value. Studies have focused on its technological and nutritional potential, highlighting the need for more research on nutrient digestibility, processing properties, and the impact of storage and processing on nutrient content.

The process of genetic enhancements in mung beans




Fig 1: The process of genetic enhancements in mung beans

The process of genetic enhancements in mung beans involves several sequential steps, starting with genetic identification. This initial phase focuses on understanding the genetic makeup of the mung beans to identify specific genes that could be modified to improve desired traits, such as yield, disease resistance, or nutritional content.

Following genetic identification, the next step is gene editing. This involves the precise modification of the identified genes. Techniques such as CRISPR-Cas9 are commonly used for this purpose, allowing scientists to add, remove, or alter genetic material at particular locations in the genome.

After modifying the genes, lab testing is conducted to assess the effects of these changes. This stage involves growing the genetically modified mung beans under controlled conditions to evaluate their growth, trait expression, and overall stability of the genetic modifications. It's a critical step to ensure that the enhancements work as intended without unwanted side effects.

The final phase is field testing, where the genetically enhanced mung beans are grown in a more natural, outdoor setting. This step is vital for assessing how the modified beans perform under varying environmental conditions, including resistance to pests and diseases, adaptability to different soil types, and the effectiveness of the genetic enhancements in real-world agricultural settings.

Each step in this process is interconnected, with the outcomes of each phase informing the next. The ultimate goal is to develop mung bean variants with improved traits that can contribute to higher yields, better nutritional value, and increased resilience to environmental stresses, thereby supporting food security and agricultural sustainability.

Genetic Diversity and Traditional Breeding

Mung bean (*Vigna radiata*) exhibits substantial genetic diversity, underpinning its adaptability to a wide range of environmental conditions and forming the basis for its breeding and improvement. Traditional breeding methods

have long leveraged this diversity to enhance yield, nutritional value, and resilience to biotic and abiotic stresses. Recent studies employing modern genetic analysis tools have provided deeper insights into the genetic diversity of mung bean, further informing breeding strategies.

Dahiya P, (2015) ^[5] explored the genetic divergence among 30 mung bean genotypes using Mahalanobis's D2 statistics, revealing six clusters indicating significant genetic diversity based on yield traits. This diversity is crucial for hybridization programs aimed at developing high-yielding varieties.

Chen *et al.* (2015) ^[3] developed and utilized microsatellite simple sequence repeat (SSR) markers to evaluate the genetic diversity and population structure of 157 mung bean accessions. The study highlighted the low polymorphism within the species, a higher polymorphic information content (PIC) for genomic SSRs (gSSRs) than for expression sequence tag SSRs (eSSRs), and distinct genetic groupings that inform breeding programs.

Kebede ZY, *et al.* (2022) ^[6] assessed genetic diversity using SSR markers in a set of mung bean accessions introduced to China from the USDA. High genetic diversity was noted, underscoring the potential of these accessions to enrich the local gene pool for breeding purposes.

Utilized RAPD profiles to determine genetic diversity among 54 mung bean accessions, including improved and local landraces. The wide range of genetic diversity observed at the molecular level corresponds to variations in morphological traits, suggesting potential targets for genetic improvement.

Molecular Breeding and Genetic Engineering

Molecular breeding and genetic engineering are transformative approaches in the field of agricultural biotechnology, significantly impacting the development of improved crop varieties, including mung beans. These techniques are vital for addressing various agricultural

challenges such as enhancing yield, improving nutritional quality, developing disease and pest resistance, and increasing tolerance to abiotic stresses like drought and salinity. The importance of these methods lies in their ability to precisely identify, manipulate, and introduce genes responsible for desirable traits, thereby facilitating the rapid and efficient improvement of crops.

Through gene discovery and characterization, scientists can uncover and understand the functions of specific genes that contribute to beneficial traits in mung beans. Marker-assisted selection speeds up the breeding process by using molecular markers to select plants that carry desired traits, making breeding more accurate and efficient. Genetic modification and genome editing technologies, such as CRISPR-Cas9, allow for the direct manipulation of the mung bean genome, enabling the addition, removal, or alteration of DNA sequences. This precision facilitates the creation of mung bean varieties with enhanced characteristics, such as increased nutritional value and resilience to environmental stresses.

Moreover, these technologies enable trait stacking, where multiple beneficial traits are combined into a single variety, optimizing the plant's performance and sustainability. By developing mung beans that require fewer agricultural inputs, such as water and pesticides, molecular breeding and genetic engineering contribute to more sustainable and environmentally friendly farming practices. Overall, these advanced techniques play a crucial role in modern agriculture, driving the progress of genetic enhancements in mung beans and ensuring food security and sustainability in the face of global challenges.

Conclusion

The analysis of genetic enhancements in mung bean reveals a focused effort on understanding and exploiting its genetic diversity for improved yield, stress resistance, and nutritional quality. Advances in genetic divergence studies, the development of molecular markers through transcriptome sequencing, and the application of genomic information for breeding underline the significant progress made in enhancing mung bean varieties. These efforts not only aim to optimize the crop's agricultural and nutritional potential but also address the challenges of biotic and abiotic stresses. As such, the ongoing research and development in mung bean genetics hold promise for achieving more resilient and nutritious mung bean cultivars, catering to the needs of both farmers and consumers globally.

References

1. Ramanujam S, Tiwari A, Mehra R. Genetic divergence and hybrid performance in mung bean. *Theoretical and Applied Genetics*,2004:45:211-214.
<https://doi.org/10.1007/BF00265001>.
2. Chen H, Qiao L, Wang L, Wang S, Blair M, Cheng X. Assessment of genetic diversity and population structure of mung bean (*Vigna radiata*) germplasm using EST-based and genomic SSR markers. *Gene*,2015:566(2):175-183.
<https://doi.org/10.1016/j.gene.2015.04.043>.
3. Chen H, Wang L, Wang S, Liu C, Blair M, Cheng X. Transcriptome Sequencing of Mung Bean (*Vigna radiata* L.) Genes and the Identification of EST-SSR Markers. *PLoS ONE*, c2015, 10.
<https://doi.org/10.1371/journal.pone.0120273>.
4. Ha J, Lee S. Mung Bean (*Vigna radiata* (L.) R. Wilczek) Breeding. In: *Advances in Plant Breeding Strategies: Legumes*, c2019.
https://doi.org/10.1007/978-3-030-23400-3_10.
5. Dahiya P, Linnemann A, Boekel M, Khetarpaul N, Grewal R, Nout M. Mung Bean: Technological and Nutritional Potential. *Critical Reviews in Food Science and Nutrition*,2015:55:670-688.
<https://doi.org/10.1080/10408398.2012.671202>.
6. Kebede ZY. Case study report on improved mung bean technology demonstration under North-West Ethiopia. *International Journal of Agriculture and Food Science*,2022:4(1):35-37.
DOI: 10.33545/2664844X.2022.v4.i1a.63