



International Journal of Multidisciplinary Research and Development



IJMIRD 2014; 1(2): 111-113
www.allsubjectjournal.com
Received: 18-07-2014
Accepted: 30-07-2014
e-ISSN: 2349-4182
p-ISSN: 2349-5979

Manoj Sharma
Department of
Mathematics RJIT, BSF
Academy, Tekanpur,

Dirichlet Average of Generalized Mittag-Leffler function and Fractional Derivative

Manoj Sharma

Abstract

In this paper Dirichlet average of a new Special function called as Generalized Mittag-Leffler function [13] which is recently given by Salim and Faraj [13]. This function has recently found essential applications in solving problems in physics, biology, engineering and applied sciences.

Keywords: Dirichlet average, Generalized Mittag-Leffler function and Fractional calculus operators.

Mathematics Subject Classification: 26A33, 33A30, 33A25 and 83C99.

1. Introduction:

Carlson [1-5] has defined Dirichlet average of functions which represents certain type of integral average with respect to Dirichlet measure. He showed that various important special functions can be derived as Dirichlet averages for the ordinary simple functions like x^t, e^x etc. He has also pointed out [3] that the hidden symmetry of all special functions which provided their various transformations can be obtained by averaging x^n, e^x etc. Thus he established a unique process towards the unification of special functions by averaging a limited number of ordinary functions. Almost all known special functions and their well known properties have been derived by this process.

In this paper the Dirichlet average of a new Special function called as Generalized M-L function has been obtained.

2. Definitions:

We give below some of the definitions which are necessary in the preparation of this paper.

2.1 Standard Simplex in $R^n, n \geq 1$:

We denote the standard simplex in $R^n, n \geq 1$ by [1, p.62].

$$E = E_n = \{S(u_1, u_2, \dots, u_n) : u_1 \geq 0, \dots, u_n \geq 0, u_1 + u_2 + \dots + u_n \leq 1\} \quad (2.1.1)$$

2.2 Dirichlet measure:

Let $b \in C^k, k \geq 2$ and let $E = E_{k-1}$ be the standard simplex in R^{k-1} . The complex measure μ_b is defined by E[1].

$$d\mu_b(u) = \frac{1}{B(b)} u_1^{b_1-1} \dots u_{k-1}^{b_{k-1}-1} (1 - u_1 - \dots - u_{k-1})^{b_k-1} du_1 \dots du_{k-1} \quad (2.2.1)$$

Will be called a Dirichlet measure.

Here

$$B(b) = B(b_1, \dots, b_k) = \frac{\Gamma(b_1) \dots \Gamma(b_k)}{\Gamma(b_1 + \dots + b_k)},$$

$$C_{>} = \{z \in \mathbb{C} : z \neq 0, |\arg z| < \pi/2\},$$

Open right half plane and $C_{>}^k$ is the k^{th} Cartesian power of $C_{>}$

2.3 Dirichlet Average [1, p.75]:

Let Ω be the convex set in $C_{>}^k$, let $z = (z_1, \dots, z_k) \in \Omega^k, k \geq 2$ and let $u.z$ be a convex combination of z_1, \dots, z_k . Let f be a measurable function on Ω and let μ_b be a Dirichlet measure on the standard simplex E in R^{k-1} . Define

Correspondence:

Manoj Sharma
Department of
Mathematics RJIT, BSF
Academy, Tekanpur,

$$F(b, z) = \int_E^0 f(u, z) d\mu_b(u) \tag{2.3.1}$$

We shall call F the Dirichlet measure of f with variables $z = (z_1, \dots, z_k)$ and parameters $b = (b_1, \dots, b_k)$. Here

$$u, z = \sum_{i=1}^k u_i z_i \text{ and } u_k = 1 - u_1 - \dots - u_{k-1} \tag{2.3.2}$$

If $k = 1$, define $F(b, z) = f(z)$.

2.4 Fractional Derivative [8, p.181]:

The concept of fractional derivative with respect to an arbitrary function has been used by Erdelyi[8]. The most common definition for the fractional derivative of order α found in the literature on the ‘‘Riemann-Liouville integral’’ is

$$D_z^\alpha F(z) = \frac{1}{\Gamma(-\alpha)} \int_0^z F(t)(z-t)^{-\alpha-1} dt \tag{2.4.1}$$

Where $Re(\alpha) < 0$ and $F(x)$ is the form of $x^p f(x)$, where $f(x)$ is analytic at $x = 0$.

3.1 Generalized Mittag-Leffler function

$$S(\beta, \beta'; x, y) = \frac{\Gamma(\beta + \beta')}{\Gamma\beta} (x - y)^{1-\beta-\beta'} D_{x-y}^{-\beta'} E_{\alpha, \beta, n}^{a_1, b_1, m}(x)(x - y)^{\beta-1} \tag{3.2}$$

Proof:

$$\begin{aligned} S(\beta, \beta'; x, y) &= \sum_{k=0}^{\infty} \frac{(a_1)_{km}}{(b_1)_{kn}} \frac{z^k}{\Gamma(\alpha k + \beta)} R_n(\beta, \beta'; x, y) \\ &= \sum_{k=0}^{\infty} \frac{(a_1)_{km}}{(b_1)_{kn}} \frac{1}{\Gamma(\alpha k + \beta)} \frac{\Gamma(\beta + \beta')}{\Gamma\beta \Gamma\beta'} \int_0^1 [ux + (1-u)y]^k u^{\beta-1} (1-u)^{\beta'-1} du \end{aligned}$$

Putting $u(x - y) = t$, we have,

$$= \sum_{k=0}^{\infty} \frac{(a_1)_{km}}{(b_1)_{kn}} \frac{1}{\Gamma(\alpha k + \beta)} \frac{\Gamma(\beta + \beta')}{\Gamma\beta \Gamma\beta'} \int_0^{x-y} [t + y]^k \left(\frac{t}{x-y}\right)^{\beta-1} \left(1 - \frac{t}{x-y}\right)^{\beta'-1} \frac{dt}{x-y}$$

On changing the order of integration and summation, we have

$$= (x - y)^{1-\beta-\beta'} \frac{\Gamma(\beta + \beta')}{\Gamma\beta \Gamma\beta'} \int_0^{x-y} \sum_{k=0}^{\infty} \frac{(a_1)_{km}}{(b_1)_{kn}} \frac{1}{\Gamma(\alpha k + \beta)} [t + y]^k (t)^{\beta-1} (x - y - t)^{\beta'-1} dt$$

Or

$$= (x - y)^{1-\beta-\beta'} \frac{\Gamma(\beta + \beta')}{\Gamma\beta \Gamma\beta'} \int_0^{x-y} E_{\alpha, \beta, n}^{a_1, b_1, m}(x) (t)^{\beta-1} (x - y - t)^{\beta'-1} dt$$

Hence, by the definition of fractional derivative, we get

$$S(\beta, \beta'; x, y) = (x - y)^{1-\beta-\beta'} \frac{\Gamma(\beta + \beta')}{\Gamma\beta} D_{x-y}^{-\beta'} E_{\alpha, \beta, n}^{a_1, b_1, m}(x)(x - y)^{\beta-1}$$

This completes the Analysis.

4. Acknowledgement:

Authors are grateful to referee for his valuable comment and improvement upon the paper.

Here , first the notation and the definition of the **Generalized Mittag-Leffler function**, introduced by Ahmad Faraj , Tariq Salim [13] has been given as

$$E_{\alpha, \beta, n}^{a_1, b_1, m}(z) = \sum_{k=0}^{\infty} \frac{(a_1)_{km}}{(b_1)_{kn}} \frac{z^k}{\Gamma(\alpha k + \beta)} \tag{1}$$

Here $\alpha, \beta \in C, Re(\alpha) > 0, Re(\beta) > 0$, $(a_1)_{km}, (b_1)_{kn}$ are the pochhammer symbols and m, n are non-negative real numbers.

3.2 Equivalence:

In this section we shall show the equivalence of single

Dirichlet average of $E_{\alpha, \beta, n}^{a_1, b_1, m}(x)$ function ($k = 2$) with the fractional derivative i.e.

References

- 3 Carlson, B.C., Special Function of Applied Mathematics, Academic Press, New York, 1977.
- 4 Carlson, B.C., Appell's function F_4 as a double average, SIAM J. Math. Anal.6 (1975), 960-965.
- 5 Carlson, B.C., Hidden symmetries of special functions, SIAM Rev. 12 (1970), 332-345.
- 6 Carlson, B.C., Dirichlet averages of $x^{-1} \log x$, SIAM J. Math. Anal. 18(2) (1987), 550-565.
- 7 Carlson, B.C., A connection between elementary functions and higher transcendental functions, SIAM J. Appl. Math. 17 (1969), 116-140.
- 8 Deora, Y. and Banerji, P.K., Double Dirichlet average of e^x using fractional derivatives, J. Fractional Calculus 3 (1993), 81-86.
- 9 Deora, Y. and Banerji, P.K., Double Dirichlet average and fractional derivatives, Rev.Tec.Ing.Univ. Zulia 16(2) (1993), 157-161.
- 10 Deora, Y, and Banerji, P,K, An Application of Fractional Calculus to the solution of Euler-Darbox equation in terma of Dirichlet average J. of fractional Calculus vol.5, may (1994) 91-94.
- 11 Erdelyi, A., Magnus, W., Oberhettinger, F. and Tricomi , F.G., Tables of Integral Transforms, Vol.2 McGraw-Hill, New York, 1954.
- 12 Gupta,S.C. and Agrawal, B.M., Dirichlet average and fractional derivatives, J. Indian Acad.Math. 12(1) (1990), 103-115.
- 13 Gupta,S.C. and Agrawal, Double Dirichlet average of e^x using fractional derivatives, Ganita Sandesh 5 (1) (1991),47-52.
- 14 Mathai. A.M. and Saxena ,R.K., The H-Function with Applications in Statistics and other Disciplines, Wiley Halsted, New York, 1978.
- 15 Salim, T, and Faraj, A ., A generalization of Mittag – Leffler function and integral operator associated with fractional calculus. J. Frac. Cal. appl. Vol.3, No.5, 1-13 (2012).
- 16 Saxena, R.K., Mathai, A. M. and Haubold, H. J., Unified fractional kinetic equation and a fractional diffusion equation, J. Astrophysics and Space Science 209 (2004) , 299-310.
- 17 Sharma, M. and Jain, R. , Dirichlet Average and Fractional Derivative, J. Indian Acad. Math.Vol.12, No. 1(1990).
- 18 Sharma, M. and Jain, R., Dirichlet Average of $\cosh x$ and Fractional Derivative, J. Indian Acad. Math.Vol.2, No. 1(2007). P17-22.